

Chemical Constituents of Peppers (*Piper* spp.) and Application to Food Preservation: Naturally Occurring Antioxidative Compounds

by Nobuji Nakatani,* Reiko Inatani,* Haruko Ohta,* and Atsuko Nishioka*

In a structure analysis of the compounds of the genus *Piper* (Family Piperaceae), we identified five phenolic amides from *Piper nigrum*, seven compounds from *P. retrofractum*, and two compounds from *P. baccatum*. All the phenolic amides possess significant antioxidant activities that are more effective than the naturally occurring antioxidant, α -tocopherol. One amide, feruperine, has antioxidant activity as high as the synthetic antioxidants, butylated hydroxyanisole (BHA) and butylated hydroxytoluene (BHT). Naturally occurring antioxidants, therefore, may surpass BHA and BHT in their ability to inactivate mutagens in food.

Introduction

Much research has involved finding ways to prevent or to delay deterioration in foods. Recently, our research (1,2) has focused on developing safe and effective compounds from natural sources (especially from edible plants) that will prolong the storage life of food.

We studied the genus *Piper* of the Family Piperaceae, to identify the structure of the compounds in its species that give organoleptic (3-5), medicinal (6-9), and insecticidal (10-13) properties. Many species, among the 700 in this genus, are used not only as spices but also as folk medicines (14). This paper reveals the structure of the compounds and their resulting properties for *Piper nigrum* L., *P. retrofractum* Vahl., and *P. baccatum* Blume.

Structural Determination of Constituents

Powdered dry fruits of white pepper (*P. nigrum*) were extracted with methylene chloride and fractionated as shown in Figure 1. The fractions determined were neutral (83.80%), weakly acidic (1.72%), strongly

acidic (0.55%), and basic (0.22%). We found piperine [1] (a pungent principle) and more than 40 constituents isolated from *P. nigrum* to be neutral compounds. The weakly acidic fraction showed significant antioxidative activity, and we subjected it to column chromatography on silica gel, using a solution of CH_2Cl_2 and MeOH as an eluant. The major compound [2a], purified upon recrystallization from hot chloroform (mp 144.0-144.5°C), produced colorless needles.

Mass spectral and elemental analyses of [2a] indicated the molecular formula $\text{C}_{18}\text{H}_{19}\text{NO}_4$.

Three absorption bands at 3500, 3360, and 3240 cm^{-1} in the infrared (IR) spectrum showed the presence of hydroxy or amino groups. The absorptions at 1645, 1615, and 980 cm^{-1} represented a *trans*-conjugated amide carbonyl group.

The λ_{max} at 295 and 321 nm in the ultraviolet (UV) spectrum indicated the presence of feruloyl moiety in the molecule, as shown in the spectrum of ferulic acid (λ_{max} 294 and 320 nm). This was also confirmed by the base peak at m/z 177 in the mass spectral (MS) data.

Further evidence for the structural characteristics was obtained from nuclear magnetic resonance (NMR) spectroscopy. One methoxy group was observed at δ 3.88, and a typical set of two doublets at δ 5.47 and δ 7.47 (both $J = 16$ Hz) was assigned to *trans*- α and *trans*- β -protons conjugated to the carbonyl group, respectively. A triplet (2H) centered at δ 2.75, ($J = 7$ Hz) and

*Department of Food and Nutrition, Faculty of Science of Living, Osaka City University, 3-3-138, Sugimoto, Sumiyoshi-ku, Osaka 558, Japan.

3. Muller, C. J., and Jennings, C. G. Constituents of black pepper. Some sesquiterpene hydrocarbons. *J. Agr. Food Chem.* 15: 762-766 (1967).
4. Muller, C. J., Creveling, R. K., and Jennings, W. G. Some minor sesquiterpene hydrocarbons of black pepper. *J. Agr. Food Chem.* 16: 113-117 (1968).
5. Traxler, J. T. Piperanine, a pungent component of black pepper. *J. Agr. Food Chem.* 19: 1135-1138 (1971).
6. Chatterjee, A., and Dutta, C. P. Alkaloids of *Piper longum* Linn.—Structure and synthesis of piperlongumine and piperlonguminine. *Tetrahedron* 23: 1769-1781 (1967).
7. Hänsel, R., Weiss, D., and Schmidt, B. Fungistatische Wirkung der Kavadooge und ihrer Inhaltsstoffe. *Planta Med.* 14: 1-9 (1966).
8. Loder, J. W., Moorhouse, A., and Russell, G. B. Tumour inhibitory plants. Amides of *Piper novae-hollandiae* (Piperaceae). *Austral. J. Chem.* 22: 1531-1538 (1969).
9. Shin, K. H., Yun, H. S., Woo, W. S., and Lee, C. K. Pharmacologically active principle of *Piper retrofractum*. *Korean J. Pharmacol.* 10: 69-71 (1979).
10. Matsui, K., Wada, K., and Munakata, K. Insect antifeeding substances in *Parabenzoin praecox* and *Piper futokadzura*. *Agr. Biol. Chem.* 40: 1045-1046 (1976).
11. Su, H. C. F. Insecticidal properties of black pepper to rice weevils and cowpea weevils. *J. Econ. Entomol.* 70: 18-21 (1977).
12. Miyakado, M., Nakayama I., Yoshioka, H., and Nakatani, N. The Piperaceae amides I: structure of piperide, a new insecticidal amide from *Piper nigrum* L. *Agr. Biol. Chem.* 43: 1609-1611 (1979).
13. Su, H. C. F., and Horvat, R. Isolation, identification, and insecticidal properties of *Piper nigrum* amides. *J. Agr. Food Chem.* 29: 115-118 (1981).
14. Atal, C. K., Dhar, K. L., and Singh, J. The chemistry of Indian *Piper* species. *Lloydia* 38: 256-264 (1974).
15. Nakatani, N., Inatani, R., and Fuwa, H. Structures and syntheses of two phenolic amides from *Piper nigrum* L. *Agr. Biol. Chem.* 44: 2831-2836 (1980).
16. Inatani, R., Nakatani, N., and Fuwa, H. Structure and synthesis of new phenolic amides from *Piper nigrum* L. *Agr. Biol. Chem.* 45: 667-673 (1981).
17. Burkill, I. H. A dictionary of the economic products of the Malay peninsula. Governments of Malaysia and Singapore, Kuala Lumpur, Malaysia, 1966, pp. 1781-1782.
18. Okogun, J. I., and Ekong, D. E. U. Extracts from the fruits of *Piper guineense* Schum. and Thonn. *J. Chem. Soc. Perkin Trans.* 1: 2195-2198 (1974).
19. Nakatani, N., and Inatani, R. Isobutyl amides from pepper (*Piper nigrum* L.). *Agr. Biol. Chem.* 45: 1473-1476 (1981).
20. Chatterjee, A., Basa, S. C., and Ray, A. B. Spectral properties of the lignan cubebin. *J. Indian Chem. Soc.* 45: 723 (1968).
21. Batterbee, J. E., Burden, R. S., Crombie, L., and Whiting, D. A. Chemistry and synthesis of the lignan (-)-cubebin. *J. Chem. Soc. (C)*: 2470-2477 (1969).
22. Branen, C. L. Toxicology and biochemistry of BHA and BHT. *J. Am. Oil Chemists Soc.* 52: 59 (1975).
23. Anon. Food Chemical News 17: 00 (March 1, 1976).
24. Sugimura, T., Nagao, M., Kawachi, T., Honda, M., Yahagi, T., Seino, Y., Sato, S., Matsukura, N., Matsushima, T., Shirai, A., Sawamura, M., and Matsumoto, H. Mutagen-carcinogens in food, with special reference to highly mutagenic pyrolytic products in broiled foods. In: *Origins of Human Cancer* (H. H. Hiatt, J. D. Watson, and J. A. Winsten, Eds.), Cold Spring Harbor Laboratory, Cold Spring Harbor, N Y, 1977, p. 1561.
25. Taylor, S. L. Mutagenesis vs. carcinogenesis. *Food Technol.* 36 (3): 65-127 (1982).
26. Pariza, M. W. Mutagens in heated foods. *Food Technol.* 36 (3): 53-56 (1982).
27. Yonei, S., and Furui, H. Lethal and mutagenic effects of malondialdehyde, a decomposition product of peroxidized lipids, on *Escherichia coli* with different DNA-repair capacities. *Mutat. Res.* 88: 23-32 (1981).
28. Shamberger, R. J., Corlett, C. L., Beaman, K. D., and Kasten, B. L. Antioxidants reduce the mutagenic effect of malonaldehyde and β -propiolactone. Part IX, Antioxidants and cancer. *Mutat. Res.* 66: 349-355 (1979).
29. Wattenberg, L. W. Inhibition of chemical carcinogen-induced pulmonary neoplasia by butylated hydroxyanisole. *J. Natl. Cancer Inst.* 50: 1541-1544 (1973).
30. McKee, R. H., and Tometsko, A. M. Inhibition of promutagen activation by the antioxidants butylated hydroxyanisole and butylated hydroxytoluene. *J. Natl. Cancer Inst.* 63: 473-477 (1979).
31. Clapp, N. K., Bowles, N. D., Satterfield, L. C., and Klima, W. C. Selective protective effect of butylated hydroxytoluene against 1,2-dimethylhydrazine carcinogenesis in BALB/c mice. *J. Natl. Cancer Inst.* 63: 1081-1085 (1979).
32. Guttenplan, J. B. Inhibition by L-ascorbate of bacterial mutagenesis induced by two N-nitroso compounds. *Nature* 268: 368-370 (1977).
33. Shamberger, R. J. Relationship of selenium to cancer. I. Inhibitory effect of selenium on carcinogenesis. *J. Natl. Cancer Inst.* 44: 931-936 (1970).
34. Sayer, J. M., Yagi, H., Wood, A. W., Conney, A. H., and Jerina, D. M. Extremely facile reaction between the ultimate carcinogen benzo[a]pyrene-7,8-diol 9,10-epoxide and ellagic acid. *J. Am. Chem. Soc.* 104: 5562-5564 (1982).